## **Dosing for Infants and Children**





From Your Healthcare Professional

#### DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

Available in:

Grape

If possible, use weight to dose; otherwise use age.

> **mL** = milliliter **tsp** = teaspoon

TYLE!	For Children
20000	ner Frant Britain
NEW	FANTS
Dasage & Directions	
Expenden Free Alcohol Free Agains Free	1
	1
Cor Only Endough Springs	Grap 1fects

### Infants' TYLENOL® Oral Suspension

Active Ingredient: Acetaminophen 160 mg (in each 5 mL)



### Children's TYLENOL® Oral Suspension

Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)

Available in:

-	Grape
4	Cherry

WEIGHT	AGE	Use only as directed.	Cherry the County as directed.
6-11 lbs	0-3 mos	1.25 mL	
12-17 lbs	4-11 mos	2.5 mL	_
18-23 lbs	12-23 mos	3.75 mL	
24-35 lbs	2-3 yrs	5 mL	<b>5 mL</b> (1 tsp)
36-47 lbs	4-5 yrs	_	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	_	10 mL (2 tsp)
60-71 lbs	9-10 yrs		<b>12.5 mL</b> (2½ tsp)
72-95 lbs	11 yrs	_	15 mL (3 tsp)

# IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: \_\_\_\_\_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Take every 4 hours as needed. Do NOT exceed more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device that comes with a specific product.
- All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).

### **Dosing for Infants and Children**





From Your Healthcare Professional

DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise use age.

> **mL** = milliliter **tsp** = teaspoon



### Infants' MOTRIN® Concentrated Drops

Active Ingredient: Ibuprofen 50 mg (NSAID)<sup>†</sup> (in each 1.25 mL)

†Nonsteroidal anti-inflammatory drug

Available in:

Dye-Free
Berry



### Children's MOTRIN® Oral Suspension

Active Ingredient: Ibuprofen 100 mg (NSAID)<sup>†</sup> (in each 5 mL or 1 tsp)

Nonsteroidal anti-inflammatory drug Use only as directed.



		Original Berry Flavor Use only as directed.  Dye-Free Berry	Use only as directed.
WEIGHT	AGE	leating Use Offity as directed.	ose only as directed.
6-11 lbs	0-5 mos	Do not use	_
12-17 lbs	6-11 mos	1.25 mL	
18-23 lbs	12-23 mos	1.875 mL	_
24-35 lbs	2-3 yrs	_	<b>5 mL</b> (1 tsp)
36-47 lbs	4-5 yrs		7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	<del>-</del>	10 mL (2 tsp)
60-71 lbs	9-10 yrs		<b>12.5 mL</b> (2½ tsp)
72-95 lbs	11 yrs	_	15 mL (3 tsp)

## IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: \_\_\_\_\_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours.
- Do NOT administer longer than 10 days, unless directed by a doctor.
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID<sup>†</sup> (prescription or non-prescription).
   MOTRIN<sup>®</sup> contains ibuprofen
- Use only the dosing device that comes with a specific product.

†Nonsteroidal anti-inflammatory drug